

Anomalous latent heat in non-equilibrium phase transitions

A.E. Allahverdyan and K.G. Petrosyan

Yerevan Physics Institute, Alikhanian Brothers St. 2, Yerevan 375036, Armenia

(Dated: February 2, 2008)

We study first-order phase transitions in a two-temperature system, where due to the time-scale separation all the basic thermodynamical quantities (free energy, entropy, etc) are well-defined. The sign of the latent heat is found to be counterintuitive: it is positive when going from the phase where the temperatures and the entropy are higher to the one where these quantities are lower. The effect exists only out of equilibrium and requires conflicting interactions. It is displayed on a lattice gas model of ferromagnetically interacting spin-1/2 particles.

PACS numbers: 64.60.Cn, 05.70.Ln, 75.10.Hk

The theory of equilibrium phase transitions is an established field with known achievements in describing transformations of various states of the matter [1, 2]. Concepts borrowed from this theory apply for some non-equilibrium transitions which get equilibrium features on the macroscopic scale, e.g., equilibrium universality classes for non-equilibrium transitions [3], transitions to the glassy state analyzed with help of effective temperature [4, 5], etc. Less is known, however, about phase transition scenarios which are impossible in equilibrium.

Here we study such a truly non-equilibrium phase transition scenario realized in the steady state of a two-temperature system. This system admits a natural generalization of the equilibrium statistical thermodynamics, i.e., the quantities like entropy, internal energy, free energy are well defined, because the constituents of the system are in local equilibrium. In spite of that, the system shows a counterintuitive type of first-order phase transition, where the latent heat is positive (*anomalous latent heat*) when transforming a high temperatures (higher entropy) phase to the low temperature (lower entropy) one, i.e., in the first phase the energy is larger. This is a non-equilibrium effect. It is well-known, even from the everyday physics, that for equilibrium transitions the latent heat is negative [1]. This plays a crucial role in the heat balance of the Earth and in the weather formation, since nearly 70% of the energy transferred from the Earth's surface is due to the latent heat consumed during the vaporization at the surface and released during the vapor condensation in the atmosphere [6].

First we shall recall the thermodynamics of two-temperature systems with different time-scales [7, 8, 9]. In contrast to the usual equilibrium case, the anomalous latent heat is not forbidden here. We then work out a simple model of mean-field Ising ferromagnet demonstrating the sought effect. A necessary condition for its existence is the presence of conflicting interactions.

Consider a pair of coupled stochastic variables s and f with Hamiltonian $H(s, f)$, which interact with different thermal baths at temperatures T_s and $T_f \equiv T$, respectively (s and f can denote a set of variables). For $T = T_s = 1/\beta$, the stationary probability distribution $P(s, f)$ of the system is Gibbsian: $P(s, f) \propto e^{-\beta H(s, f)}$. For $T \neq T_s$ we can derive the stationary $P(s, f)$ if the

variables have different characteristic times: s is slow, while f is fast (adiabatic limit). This derivation together with corrections coming from a large, but finite time-scale difference was given in [8] based on stochastic equations of motion. Here we recall the heuristics [7, 8, 9, 10]. On the times relevant for f , s is fixed, and the conditional probability $P(f|s)$ is Gibbsian:

$$P(f|s) = \frac{1}{Z(s)} e^{-\beta H(s, f)}, \quad Z(s) = \text{Tr}_f e^{-\beta H(s, f)} \quad (1)$$

where Tr_f is the sum over all values of f . The steady-state $P(s)$ is found by noting that on the times relevant for s , τ is already in the conditional steady state. Thus the force $\partial_s H(s, f)$ acting on s can be averaged over $P(f|s)$: $\text{Tr}_f [\partial_s H(s, f) P(f|s)] = \partial_s F(s)$, where $F(s) = -T \ln Z(s)$ is the conditional free energy. The steady $P(s)$ is Gibbsian with the Hamiltonian $F(s)$:

$$P(s) = \frac{e^{-\beta_s F(s)}}{\mathcal{Z}} = \frac{Z_{T_s}^T(s)}{\mathcal{Z}}, \quad \mathcal{Z} = \text{Tr}_s e^{-\beta_s F(s)}, \quad (2)$$

and the common probability is $P(s, f) = P(s)P(f|s)$.

This two-temperature situation admits a (generalized) thermodynamical description, because in the adiabatic limit both variables are in local equilibrium [7, 8, 9]. (It has certain analogies with non-equilibrium thermodynamics of the glassy state proposed in [5].) The average energy of the system is $\mathcal{U} = \text{Tr}_{s, f} [P(s, f) H(s, f)]$, or

$$\mathcal{U} = -\frac{1}{n} \partial_\beta \ln \mathcal{Z}|_n, \quad n \equiv \frac{T}{T_s} \quad (3)$$

where the derivative is taken for fixed n and we used Eqs.(1, 2). $\mathcal{Z}$ is a generating function, since an arbitrary average is found via the proper source term analogously to (2). The entropies of s and f are, respectively,

$$\mathcal{S}_s = -\text{Tr}_s P(s) \ln P(s), \quad (4)$$

$$\mathcal{S} = -\text{Tr}_s P(s) [\text{Tr}_f P(f|s) \ln P(s|f)]. \quad (5)$$

Eq.(4) is the usual definition of entropy. Eq.(5) is the conditional entropy; it appears due to the adiabatic limit. The total entropy is the sum of partial ones $\mathcal{S}_{\text{tot}} \equiv -\text{Tr}_{s, f} [P(s, f) \ln P(s, f)] = \mathcal{S} + \mathcal{S}_s$. Now the

steady distributions (1, 2) are obtained when minimizing the free energy of the slow motion

$$\mathcal{F} = -T_s \ln \mathcal{Z}. \quad (6)$$

To this end, note from (1–5) that the free energy and the entropies are expressed as

$$\mathcal{F} = \mathcal{U} - T\mathcal{S} - T_s\mathcal{S}_s, \quad (7)$$

$$\mathcal{S}_s = -\partial_{T_s}\mathcal{F}|_T, \quad \mathcal{S} = -\partial_T\mathcal{F}|_{T_s}, \quad (8)$$

generalizing the usual thermodynamical relations.

Let us now study the first-order phase transitions in this approach. Consider two phases l and h . Assume that when T_s decreases below some critical value $T_s^{(c)}$ (T is fixed), the phase l dominates, since its free energy is smaller: $\mathcal{F}_l < \mathcal{F}_h$. For $T_s > T_s^{(c)}$ the dominating phase is h : $\mathcal{F}_l > \mathcal{F}_h$. At high (low) temperatures T_s , l (h) is metastable. $\mathcal{F}_l = \mathcal{F}_h$ at $T_s = T_s^{(c)}$, while other quantities change by jump. Denoting $\delta X = X_l - X_h$ we get

$$\delta\mathcal{U} = T_s\delta\mathcal{S}_s + T\delta\mathcal{S} = -T_s\partial_{T_s}[\delta\mathcal{F}]|_T - T\partial_T[\delta\mathcal{F}]|_{T_s}, \quad (9)$$

where $\delta\mathcal{U}$ is the latent heat of the transition. Note that in general both $\delta\mathcal{S}_s$ and $\delta\mathcal{S}$ are functions of T_s and T . In the vicinity of $T_s^{(c)}$, $\delta\mathcal{F} = \mathcal{F}_l - \mathcal{F}_h$ is an increasing function of T_s (for fixed T); thus $\delta\mathcal{S}_s < 0$. The sign of $\delta\mathcal{S}$ is, however, left open and we cannot conclude that $\delta\mathcal{U} < 0$. It may be even positive (anomalous latent heat), provided $\delta\mathcal{S} > 0$, that is provided the system moves from l to h when decreasing T for constant T_s .

Things get different if we decrease both T_s and T , with constant $n = T/T_s$. Since $\delta\mathcal{F}(T_s, nT_s) = \mathcal{F}_l - \mathcal{F}_h$ has to be an increasing function of T_s , we get $\frac{d}{dT_s}\delta\mathcal{F} = (\partial_{T_s} + n\partial_T)\delta\mathcal{F} > 0$, and then (9) shows that $\delta\mathcal{U} < 0$ confirming the equilibrium result for $T_s = T$ [1].

Thus for $T_s \neq T$ the anomalous latent heat is not forbidden, provided the transition is not driven by a proportional change of both temperatures. Here is a model demonstrating this effect. Particles are located at the nodes of a lattice embedded into a particle reservoir with chemical potential $\mu = -\alpha < 0$. Each particle carries an Ising spin. At short distances the particles repel each other so that not more than one particle can occupy a single node. The Ising spins interact ferromagnetically. This interaction is active between two nodes only if they both are occupied; $s \equiv \{s_i = \pm 1\}_{i=1}^N$ and $f \equiv \{f_i = 0, 1\}_{i=1}^N$ are, respectively, the spins and the occupations of N nodes. We assume all nodes can interact with each other (mean-field). The Hamiltonian reads

$$H(s, f) = -\frac{J}{2N} \sum_{i \neq k}^N f_i f_k s_i s_k + \alpha \sum_i f_i, \quad (10)$$

where $J/(2N) > 0$ is the ferromagnetic coupling constant with $J = \mathcal{O}(1)$, as required for the extensivity.

We make two assumptions. 1) s and f couple, respectively, to a spin bath and lattice bath at different temperatures, $T_s = \frac{1}{\beta_s}$ and $T = \frac{1}{\beta}$. 2) s (f) is slow

(fast), i.e., the relaxation time τ_f of f is much shorter than the relaxation time τ_s of s . These times are driven by the interactions with thermal baths, since H contains only commuting terms; note that in the quantum setting $s_i = s_{i,z}$ and $f_i = \frac{1}{2}(1 + \sigma_{i,z})$, where $s_{i,z}$ and $\sigma_{i,z}$ are the third Pauli matrices.

The above assumptions are motivated by NMR/ESR physics, where for nuclear or electronic spins the spin bath is realized by relatively weak dipole interactions and does indeed lead to a temperature different from the lattice one [11]. This spin temperature can be tuned by external fields or by spin cooling and plays an important role in analyzing experiments [11]. The relaxation time τ_s on which the spin temperature is established is known as T_2 time in NMR/ESR, and usually varies between 10^{-4} s and 1s depending on the material. The spin and the lattice temperatures tend to equalize on the T_1 time, which for many magnetic materials amounts to minutes or hours [11]. Estimating the relaxation time τ_f of f via the rotation-vibration mechanism as $10^{-12} - 10^{-9}$ s [11], we see that our assumptions are based on $\tau_f \ll \tau_s \ll T_1$.

The model is a suitable candidate for displaying the anomalous latent heat, since the two terms in the RHS of (10) are in conflict: for two lined up spins, $s_i s_k > 0$, the term $-\gamma \sum_{\langle i,k \rangle} f_i f_k s_i s_k$ tends to increase occupations, $f_i f_k > 0$, while $\alpha \sum_i f_i$ makes this increase costly. The equilibrium ($T_s = T$) version of the model is related to Blume-Capel model [2]. Similar models were employed recently for describing solid mixtures [2], inverse freezing (reentrance) phenomena [12] and glassy physics [13].

Employing $e^{(\sum_i f_i s_i)^2} = c \int dm e^{-\frac{N}{2}\beta J m^2 + \beta J m \sum_i s_i f_i}$, where $c = \sqrt{\frac{N\beta J}{2\pi}}$, we express from Eqs.(2, 10) the partition function $\mathcal{Z}$ of the model as

$$\mathcal{Z} = \text{Tr}_s \left[\text{Tr}_f e^{\frac{\beta J}{2N} (\sum_i f_i s_i)^2 - \alpha \beta \sum_i f_i} \right]^n = \int \mathcal{D}m e^{-N\beta_s \mathcal{F}},$$

where $\mathcal{D}m \equiv c^{\frac{n}{2}} \prod_{a=1}^n \int dm_a$, and we assumed that $n \equiv \frac{T}{T_s}$ is an integer (replica method); later on we shall make continuation to real n . We also defined $\phi(m) = 1 + e^{-\alpha\beta + \beta J m}$ and

$$\beta_s \mathcal{F}(m) = \frac{\beta J}{2} \sum_{a=1}^n m_a^2 - \ln \left[\sum_{k=\pm 1} e^{\sum_{a=1}^n \ln \phi(k m_a)} \right]. \quad (11)$$

$\mathcal{Z}$ is calculated by the saddle-point method, where one searches the deepest minimum of $\mathcal{F}$ as a function of m_a . This minimum is reached for $m_a = m$ (replica symmetry), where m satisfies the stationarity condition $\frac{\partial}{\partial m_a} \mathcal{F}|_{m_a=m} = 0$, i.e.,

$$m = e^{-\alpha\beta} \frac{\phi^{n-1}(m) e^{\beta J m} - \phi^{n-1}(-m) e^{-\beta J m}}{\phi^n(m) + \phi^n(-m)}. \quad (12)$$

The corresponding value of $\mathcal{F}$ obtained from (11) is

$$\mathcal{F} = \frac{Jm^2}{2} - T_s \ln [\phi^n(m) + \phi^n(-m)]. \quad (13)$$

It is seen that $\mathcal{F}$ is the free energy of the system and that m is the magnetization:

$$m = \frac{1}{N} \sum_{i=1}^N \langle \tau_i \sigma_i \rangle. \quad (14)$$

Due to the symmetry of the model, Eq.(12) admits two equivalent solutions with $\pm m$. The choice between them is done spontaneously.

The energy $\mathcal{U}$ and the entropies of the spins $\mathcal{S}_s$ and the occupations $\mathcal{S}$ are calculated from Eqs.(3-10)

$$\mathcal{U} = -\frac{Jm^2}{2} + \frac{\alpha}{z(m)} e^{-\alpha\beta} \sum_{k=\pm 1} \phi^{n-1}(km) e^{\beta Jkm}, \quad (15)$$

$$\mathcal{S}_s = \ln z(m) - \frac{n}{z(m)} \sum_{k=\pm 1} \phi^n(km) \ln \phi(km), \quad (16)$$

$$\mathcal{S} = \beta \mathcal{U} + \frac{1}{z(m)} \sum_{k=\pm 1} \phi^n(km) \ln \phi(km) - \frac{\beta Jm^2}{2}, \quad (17)$$

where $z(m) = \sum_{k=\pm 1} \phi^n(km)$. The two terms in RHS of (15) are equal, respectively, to the spin-spin interaction energy $\frac{J}{2N} \langle (\sum_i f_i s_i)^2 \rangle$ and $\alpha \mathcal{N}$, where $\mathcal{N} = \sum_i \langle f_i \rangle$ is the average number of particles.

In the derivation of (15–17) we assumed that the system is in the ferromagnetic phase: $m \neq 0$. For the paramagnetic phase $m = 0$ one has from (13)

$$\mathcal{U} = \frac{\alpha}{e^{\alpha\beta} + 1}, \quad \mathcal{S}_s = \ln 2, \quad \mathcal{S} = \frac{\alpha\beta}{e^{\alpha\beta} + 1} + \ln(1 + e^{-\alpha\beta}).$$

The phase diagram of the model is constructed in terms of three adimensional parameters:

$$a \equiv \alpha/J, \quad \theta_s = T_s/J, \quad \theta = T/J. \quad (18)$$

Let us start with few particular cases. For $\theta \rightarrow 0$ we get from (13) (assuming $m \geq 0$)

$$\beta_s \mathcal{F} = \frac{m^2}{2\theta_s} - \ln[1 + \zeta(m-a) + \zeta(a-m)e^{(m-a)/\theta_s}], \quad (19)$$

$$m = (1 + e^{(a-m)/\theta_s})^{-1} (1 - \zeta(a-m)), \quad (20)$$

where $\zeta(x)$ is approximated by the step function: $\zeta(x) = 1(0)$ for $x > 0(x < 0)$. We see that $m \neq 0$ only for $a < 1$. However, only for $a < 0.5$, Eq. (20) predicts first-order phase transition to the ferromagnet at $\theta_s = \theta_s^{(c)}$. Above this temperature there are no particles in the system, $\mathcal{N} = 0$, since the energy cost for consuming a particle from the particle reservoir is too high (due to $T \rightarrow 0$). At $\theta_s = \theta_s^{(c)}$ a finite fraction of particles is consumed from the particle reservoir making up the ferromagnetic phase, which appears as a metastable state at a higher critical temperature $\theta_s^* > \theta_s^{(c)}$. There is no first-order transition for $a > 0.5$, though for $1 > a > 0.5$ the ferromagnet can exist as a metastable phase.

In a different limit $\alpha = 0$ we get from (13):

$$\frac{\mathcal{F}}{J} = \frac{m^2}{2} - \theta_s \ln \cosh\left(\frac{m}{2\theta_s}\right) - \theta \ln \cosh\left(\frac{m}{2\theta}\right), \quad (21)$$

$$2m = \tanh\left(\frac{m}{2\theta}\right) + \tanh\left(\frac{m}{2\theta_s}\right), \quad (22)$$




FIG. 1: The phase diagram on the $T-T_s$ plane; $\frac{\alpha}{J} = 0.45$. **P** and **F** refer to the paramagnet and ferromagnet, respectively. Bold line: first-order phase transitions. Normal line: second-order phase transitions. For $T \rightarrow \infty$ this line monotonically saturates at $T_s = 0.25J$, as seen from (23). Dashed line: the instability line of the paramagnet. On the AC segment the latent heat is anomalous.

Now there are only second-order phase transitions for the same reason as for the usual (Curie-Weiss) mean-field ferromagnet: $\partial_m^3 F$ can never turn to zero for $m \neq 0$. The phase diagram on the $\theta_s - \theta$ plane amounts to the ferromagnet (paramagnet) located below (above) the line

$$\theta^{-1} + \theta_s^{-1} = 4. \quad (23)$$

The same Eqs.(21, 22) without the terms containing θ apply for $\theta \rightarrow \infty$. Here f_i fluctuate so strongly that the influence of α disappears.

A necessary condition for second-order transition is the local instability of the paramagnet: $\partial_m^2 \mathcal{F}|_{m=0} = 0$, i.e.,

$$\theta_s = [1 + e^{a/\theta} (2 + e^{a/\theta} - \theta^{-1})]^{-1}. \quad (24)$$

We could continue this reasoning and develop the Ginzburg-Landau expansion, but here it is easier to study (12, 13) directly. For $a > 0.5$ there is only the second-order phase transition to ferromagnet at temperatures given by (24). Eq.(14) shows that m can increase by lining up the spins with or without increasing the occupations. For $a > 0.5$ the second way works, since it is too costly to absorb particles from the reservoir. Indeed, slightly below the transition the difference in the average number of particles is small: $\Delta \mathcal{N} \propto m^2 \ll m$; see (15).

For $0.38 < a < 0.5$ the phase diagram is of the type presented in Fig.1. Now there are first-order phase transitions from the paramagnet to the ferromagnet. Recall Eqs.(19, 20) for understanding their mechanism. The transition (phase coexistence) line is found by solving (14), looking for the deepest minima of $\mathcal{F}(m)$, and requiring the continuous change of the free energy: $\mathcal{F}(m) = \mathcal{F}(0)$. The ferromagnet corresponds to $\mathcal{F}(m) < \mathcal{F}(0)$ (recall the saddle-point method.) Other quantities of interest change by jump. The ferromagnet first appears as a metastable phase above the first-order transition line, while the paramagnet survives as metastable till the dashed line on Fig.1. Thus, in the vicinity of the first-order transition line, both phases are locally stable. This




FIG. 2: The same as in Fig.(1), but now $\alpha = 0.245 J$.

bistability implies hysteresis and memory: when changing the temperature the final state of the system (para or ferromagnet) depends on its initial state. Since both m and $\mathcal{N}$ increase in the ferromagnet, m jumps to a non-zero value not simply due to lining up the existing spins, but also due to absorption of additional particles from the reservoir.

On the AC segment the system moves from the paramagnet to the ferromagnet upon *increasing* T or *decreasing* T_s . Thus, $\mathcal{S}_s$ is smaller in the ferromagnet, while $\mathcal{S}$ is larger there: $\delta\mathcal{S}_s < 0$ and $\delta\mathcal{S} > 0$; see (9). This is necessary for the existence of the anomalous latent heat effect; see (9) and below. On the AB segment the latent heat is *anomalous*: when decreasing T_s for a fixed T the system moves towards the ferromagnet and its internal energy *increases*. This is due to a positive energy brought about by the particles coming from the reservoir; see (15). The maximal magnitude of the effect reached in the middle of AC segment is $\delta\mathcal{U} \sim J/20$. Recall that the transition to the ferromagnet is induced by the tendency of the spin interaction energy to decrease. The latent $\delta\mathcal{U}$ is a consequence of this tendency that for $\delta\mathcal{U} > 0$ overcomes its cause. On the whole first-order transition line the total entropy $\mathcal{S}_{\text{tot}} = \mathcal{S} + \mathcal{S}_s$, given by (16, 17), jumps by a negative amount: expectedly the ferromagnet is more ordered than the paramagnet.

At the point B the latent heat is zero, while the BC segment shows an anomalous latent heat in a different

scenario: upon *decreasing* T the system goes to the paramagnet, where the energy is *higher*. The main difference between the two scenarios is that now the low- T phase (paramagnet) has higher total entropy $\mathcal{S}_{\text{tot}}$: a first-order transition to a higher entropy phase is induced by decreasing the temperature T . This is impossible in equilibrium; see Eq.(9).

An example of the phase diagram for $a < 0.38$ is presented in Fig.2. The qualitative shape of the ferromagnet-paramagnet boundary is given by (23). The transitions to the ferromagnet are always induced when decreasing either temperature: there is no anomalous latent heat here. During the first-order transitions the particle number still increases: $\delta\mathcal{N} > 0$, but now this brings to decreasing entropy $\delta\mathcal{S} < 0$. In the present low- a regime the occupations are slaved by the spins so that the incoming particles arrange their entropy in the way dictated by the spin-spin interaction energy. For $\frac{a}{J} \rightarrow 0$, the first-order transitions gradually disappear.

In conclusion, we found first-order phase-transitions with an unexpected sign of the latent heat: it is positive when going from high temperature (high entropy) phase to the low temperature (low entropy) one. Two conditions are necessary for this: global non-equilibrium (two different temperatures), though the local equilibrium is kept and leads to a slightly generalized thermodynamics, and conflicting interactions. The effect was displayed on the model of lattice gas whose particles carry ferromagnetically interacting Ising spins. Motivating by NMR/ESR physics we assumed that the spins equilibrate at a temperature different from the lattice one. For this model there is a conflict between attractive interaction facilitating the income of particles from the reservoir and the chemical potential that makes this income costly.

The effect seems to be generic, since we found it in many other situations, e.g., in the present model, but with fast spins and slow occupations, models with multi-spin interaction, the Ising spin-link ferromagnet, etc.

Thanks to Theo Nieuwenhuizen for discussions.

A.E. A. was supported by CRDF grant ARP2-2647-YE-05. K.G. P. was supported by ISTC grant A-820.

-
- [1] L.D. Landau and E.M. Lifshitz, *Statistical Physics, I* (Pergamon Press Oxford, 1978).
 - [2] D.A. Lavis and G.M. Bell, *Statistical Mechanics of Lattice Systems*, vol. 1 (Springer-Verlag, 1999).
 - [3] B. Schmittmann and R.K.P. Zia, in *Phase Transitions and Critical Phenomena*, eds. C. Domb and J.L. Lebowitz (Academic Press, NY, 1996).
 - [4] L.F. Cugliandolo *et al.*, Phys. Rev. E, **55**, 3898 (1997).
 - [5] Th.M. Nieuwenhuizen, Phys. Rev. Lett., **80**, 5581 (1998); Phys. Rev. E **61**, 267 (2000).
 - [6] D.L. Hartmann, *Global Physical Climatology* (San Diego, CA, Academic Press 1994).
 - [7] R. Landauer and J. Woo, Phys. Rev. A **6**, 2205 (1972).
 - [8] A.E. Allahverdyan and Th.M. Nieuwenhuizen, Phys. Rev. E, **62**, 845 (2000).
 - [9] O.M. Ritter *et al.* Phys. Rev. E **69**, 016119 (2004).
 - [10] A.C.C. Coolen *et al.*, J. Phys. A **26**, 3681 (1993). A.E. Allahverdyan *et al.*, Eur. Phys. J. B, **16**, 317, (2000).
 - [11] M. Goldman, *Spin Temperature and NMR in Solids* (Clarendon Press, Oxford, 1970).
 - [12] N. Schupper and N.M. Shnerb, Phys. Rev. Lett., **93**, 037202 (2004).
 - [13] A. Crisanti and L. Leuzzi, Phys. Rev. Lett., **89**, 237204 (2002). M. Selitto and J. Kurchan, cond-mat/0507128.